



# Immunization Timing

Age 2 months	Interval from previous dose
DTaP	
Hib	
PCV (Pneumo)	
Polio (IPV)	
RV <sup>2</sup> (Rotavirus)	
HepB <sup>1</sup>	

Age 4 months	Interval from previous dose
DTaP	1-2 months
Hib	1-2 months
PCV (Pneumo)	1-2 months
Polio (IPV)	1-2 months
RV <sup>2</sup> (Rotavirus)	4-10 weeks
HepB <sup>1</sup>	

Age 6 months	Interval from previous dose
DTaP	1-2 months
Hib <sup>3</sup>	1-2 months
PCV (Pneumo)	1-2 months
Polio <sup>6</sup> (IPV)	1-14 months
RV <sup>2</sup> (Rotavirus)	4-10 weeks
HepB <sup>1</sup>	

Age 12 months	Interval from previous dose
DTaP <sup>4</sup>	
Hib <sup>4</sup>	2-9 months
PCV <sup>4</sup> (Pneumo)	2-6 months
MMR <sup>5</sup>	
Varicella <sup>5</sup> (Chickenpox)	
HepA	

Age 18 months	Interval from previous dose
HepA	6-12 months

Age 6 mos and older	and high risk kids
Flu	Every Year

**Early Childhood Program Entry (Age Appropriate)**

DTaP, Polio, Hib, Pneumococcal, Hepatitis A, Hepatitis B, MMR, Varicella

---

**Kindergarten Entry**

DTaP, Polio, MMR, Varicella, Hepatitis A, Hepatitis B



Reproduced with permission from the California Department of Health Services, Immunization Branch.

1-800-275-0659 • [www.immunize-utah.org](http://www.immunize-utah.org)

This is a suggested schedule. Your child's immunization schedule may vary depending upon your health care provider's judgment. For more information, consult with your health care provider and/or the "Recommended Childhood Immunization Schedule, United States, 2010" at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).

- <sup>1</sup> Hepatitis B series: The Hepatitis B vaccine is a three dose series. If Hepatitis B vaccine is not given at birth, the first dose can be given at 2 months of age. The second dose should be given at 4 months of age. The third dose should be given at  $\geq 24$  weeks of age. If the combination vaccine ComVax<sup>®</sup> (combined Hib and Hepatitis B) is used, a fourth dose can be given. When using Comvax<sup>®</sup>, the final dose is given at 12-15 months of age.
- <sup>2</sup> RotaTeq<sup>®</sup> vaccine is a three-dose series given at 2, 4, and 6 months of age. Rotarix<sup>®</sup> vaccine is a two-dose series given at 2 and 4 months of age. Vaccination with either rotavirus vaccine should not be started for infants 15 weeks of age or older. All doses should be completed by 8 months of age.
- <sup>3</sup> The number of doses in the Hib series depends on the type of vaccine used. This six month dose of Hib vaccine can be skipped if certain combination vaccines (PedVax<sup>®</sup> and/or ComVax<sup>®</sup>) are used.
- <sup>4</sup> The final dose of DTaP, Hib and Pneumococcal (PCV) vaccines should be given  $\geq 12$  months of age. For school entry, one dose of DTaP should be given  $\geq 4$  years of age.
- <sup>5</sup> The combination vaccine, MMRV (Measles, Mumps, Rubella, Varicella), can be given when both the MMR and the Varicella (Chickenpox) vaccines are necessary.
- <sup>6</sup> If Pentacel<sup>®</sup> (DTaP, IPV, Hib) vaccine is used, a fourth dose of IPV at 12-15 months of age may be given. For school entry, one dose of IPV should be given  $\geq 4$  years of age.